

PRODUCT INFORMATION SHEET ZINPRO® 40

A nutritional feed ingredient for animal feed. When used as a commercial feed ingredient, it must be declared as zinc methionine complex. Association of American Feed Control Officials (AAFCO) No. 57.151.



TYPICAL ANALYSIS

MINERALS	Zinc	4.0%
PROXIMATES	Moisture	1.5 - 3.0%
	Ash	40.0%
	Crude Protein	6.0%

PHYSICAL CHARACTERISTICS

Color	Brown		
Texture	Granular		
Particle Characteristics	Free flowing, slightly hygroscopic		
Density	610 - 670 kg/m ³		
Solubility	Active product is water soluble		
Particle Size	150 μm < 90% < 850 μm		
Package Type	Multiwall bag		
Package Weight	25 kg		

SAFE HANDLING & STORAGE

When correctly used, there is no toxicity hazard in the use of ZINPRO[®] 40. Refer to the SDS for more detail. Store in a clean and dry environment. Follow label directions and rotate inventory to ensure fresh product.

FEEDING INSTRUCTIONS

		g/hd/d	g/MT	PPM (mineral)*
DAIRY CATTLE	All			40
BEEF CATTLE	Creep Feed		1,610	
	Weaned Calves			30
	Cows and Bulls	9		
	Feedlot Receiving			60
	Feedlot Finishing			60
SHEEP	All†	1		
GOATS	All†	1.25		
HORSES	All	10	2,500	
SWINE	Nursery [†]		1,250	
	Grow/Finish [†]		1,250	
	Sows and Boars [†]		1,250	
POULTRY	All		1,000	
AQUACULTURE	Salmonids and Marine Fish		2,000	
	Tilapia and Freshwater Fish		1,500	
	Shrimp		1,500	
COMPANION	Canine		2,500	
	Feline		1,875	
SPECIALTY	Deer [†]	4.5		
	Elk†	9		
	Llamas [†]	2.5		
	Rabbits [†]	0.28		

* PPM of mineral provided in the complete diet.

† Double recommended levels during periods of reduced feed intake, or high production stress.

BEEF CATTLE: Creep feed recommendation based on daily intake of 1% BW. HORSES: All recommendations based on a mature horse of 500 kg BW consuming 4 kg of complete feed.